

On Initial Coefficient Inequalities for New Subclasses of Meromorphic Bi-univalent Functions

F. Müge Sakar

Batman University, Faculty of Management and Economics, Department of Business Administration, Batman-TURKEY
mugesakar@hotmail.com

Abstract

In this present study, we investigate two new subclasses of meromorphic and bi-univalent functions defined on $\Delta = \{z : z \in \mathbb{C}, 1 < |z| < \infty\}$. We obtain improved estimates on the initial Taylor-Maclaurin coefficients for the functions in these subclasses. Some other closely related results are also indicated.

References

1. P.L. DUREN. Coefficient of Meromorphic Schlicht Functions. Proc. Amer. Math. Soc. 28 (1971) 169-172.
2. D. A. BRANNAN AND J. CLUNIE. Aspects of Contemporary Complex Analysis. in:Proceedings of the NATO Advanced Study Institute held in Durham, England, August 26 September 6, 1974, Academic Press, New York, NY, USA, 1979.
3. J. CLUNIE AND F. R. KEOGH. On Starlike and Convex Schlicht Functions. MR 22, 1682, J. Lond. math. Soc. 35 (1960) 229-233.
4. H.M. SRIVASTAVA AND A.K. MISHRA AND P. GOCHHAYAT. Certain Subclasses of Analytic and Bi-univalent Functions. Appl. Math. Lett. 23 (10) (2010) 1188-1192.
5. G. SPRINGER. The Coefficient Problem for Schlicht Mappings of the Exterior of the Unit Circle. Amer. Math. Soc. 70 (1951) 421-450.